

**Universitat de Lleida**

Document downloaded from:

<http://hdl.handle.net/10459.1/72362>

The final publication is available at:

<https://doi.org/10.1007/s13355-017-0543-7>

Copyright

(c) The Japanese Society of Applied Entomology and Zoology, 2018

[Click here to view linked References](#)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 1 Title: Captures of oriental fruit moth, *Grapholita molesta* (Lepidoptera: Tortricidae), in  
12  
13 2 traps baited with host-plant volatiles in Chile  
14  
15 3 Authors: Wilson Barros-Parada<sup>1\*</sup>, Byrappa Ammagarahalli<sup>2‡</sup>, Esteban Basoalto<sup>3</sup>, Eduardo  
16  
17 4 Fuentes-Contreras<sup>1</sup> and César Gemenio<sup>2§</sup>  
18  
19 5  
20  
21 6 <sup>1</sup> Millennium Nucleus Center in Molecular Ecology and Evolutionary Applications in the  
22  
23 7 Agroecosystems (CEM), Facultad de Ciencias Agrarias, Universidad de Talca, Casilla 747,  
24  
25 8 Talca, Chile; e-mails: wbarros@utalca.cl, efuentes@utalca.cl  
26  
27 9 <sup>2</sup> Department of Crop and Forest Sciences, University of Lleida, Av. Alcalde Rovira Roure  
28  
29 10 191, 25198 Lleida, Spain; e-mails: nnnbyrareddy20@gmail.com,  
30  
31 11 cesar.gemenio@pvcf.udl.cat  
32  
33 12 <sup>3</sup> Instituto de Producción y Sanidad Vegetal, Facultad de Ciencias Agrarias, Universidad  
34  
35 13 Austral de Chile, Casilla 567, Valdivia, Chile; e-mail: esteban.basoalto@uach.cl  
36  
37 14  
38  
39 15 \* Share equal credit  
40  
41 16 ¶ Present address: Escuela de Agronomía, Pontificia Universidad Católica de Valparaíso,  
42  
43 17 Casilla 4-D, Quillota, Chile.  
44  
45 18 ‡ Present address: Department of Biological Sciences, University of Cincinnati, Cincinnati,  
46  
47 19 Ohio 45221, USA.  
48  
49 20 § Corresponding author, e-mail: cesar.gemenio@pvcf.udl.cat  
50  
51 21  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Formatted: Spanish (Spain)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

22 Abstract

23

24 Studies in Australia and China identified host-plant volatile blends from peach and pear  
25 that captured relatively high numbers of *Grapholita molesta* (Busck). To determine if  
26 these blends are attractants in other countries and relative to each other, the two host-  
27 plant blends, a laboratory blend identified in Switzerland, and a new "Total blend" made  
28 by mixing components of all three blends, were field tested in Chile for the first time.  
29 Same solvent type, concentrations and dispensers as in the original studies, plus an  
30 additional concentration and solvent, were used. Only the Swiss blend at the low *n*-  
31 hexane concentration captured significantly more males than the solvent traps, yet, in  
32 very low numbers ( $1.46 \pm 1.46$ , mean  $\pm$  SEM males/trap/week). Furthermore, host-plant  
33 blends decreased male captures in sex pheromone traps, and the effect was dose-  
34 dependent for the Chinese and Total blends. A laboratory flight-tunnel test confirmed the  
35 lack of *G. molesta* male response to the Australian, Chinese and Swiss plant blends. In the  
36 flight tunnel, however, the males responded sooner and in higher numbers to mixtures of  
37 sex pheromone with host-plant blends than they did to the sex pheromone alone.

38

39 Keywords

40 host-plant volatiles, sex pheromone, synergism, flight tunnel, traps

41

42 Introduction

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

43 Moths rely on their sense of smell to locate mates. Females release relatively small  
44 quantities of highly volatile pheromone molecules detected by very sensitive receptors on  
45 the male antennae (Allison and Cardé 2016). The high sensitivity and species-specificity of  
46 moth sex pheromones and their strong effect on males has made them a cornerstone tool  
47 in moth pest control. Pheromones are used to monitor insect-pest occurrence, time  
48 insecticide applications, bring the insect in contact with insecticides, or remove a  
49 significant part of the population. Foremost, sex pheromones are used to disrupt mating  
50 and so reduce population levels and crop damage (Miller and Gut 2015; Witzgall et al.  
51 2010). One way to determine if mating disruption negatively affects male attraction is to  
52 use sex pheromone traps to compare male captures in pheromone-disrupted and non-  
53 disrupted crops. However, not catching males with pheromone traps does not necessarily  
54 imply a failure of males to find females (Knight et al. 2013). In addition, pheromone traps  
55 attract only males, thus, no information is available on the mating status of females (Light  
56 et al. 2017). One method to facilitate monitoring of males and females in sex pheromone  
57 disrupted crops is to use host-plant volatile lures (Miller and Gut 2015). Despite the  
58 potential importance of plant volatiles in pest management, there are relatively few  
59 commercial plant volatile attractants for moth control (Szendrei and Rodriguez-Saona  
60 2010). One of them is the pear ester, ethyl (*E,Z*)-2,4-decadienoate, a volatile from ripe  
61 pears that is a relatively selective attractant of male and female *Cydia pomonella* (L.) and  
62 is commercialized as a combo lure together with the pheromone for monitoring and  
63 increasing the efficiency of mating disruption (Knight et al. 2014; Light et al. 2017).

64 The oriental fruit moth, *Grapholita molesta* (Busck) is a worldwide pest of peach  
65 (*Prunus persica* (L.)), apple (*Malus domestica* Borkh), and other stone and pome fruit tree  
66 species (Rothschild and Vickers 1991). Early in the season the larva bores on green shoots,  
67 moving to new ones as they are consumed. This feeding hinders the formation of new  
68 branches and causes problems in tree nurseries. Later in the season, when shoots start to  
69 harden, feeding shifts to newly available fruit, which can cause major economic loss. *G.*  
70 *molesta* is primarily a pest of peach, but in recent years there have been increasing  
71 reports of damage to apple fruit worldwide (Wei et al. 2015). The selection of oviposition  
72 locations is thus vital and is aided, at least in part, by host-plant volatiles (Myers et al.,  
73 2007; Piñero and Dorn 2007).

74 The control of *G. molesta* relies strongly on repeated insecticide applications  
75 throughout the season, with the well-known negative impacts on humans and the  
76 environment (Guillette and Iguchi 2012). The release of the synthetic pheromone blends  
77 from passive dispensers or puffers results in mating disruption and population control  
78 (King et al. 2014, Witzgall et al. 2010). Although there are no commercial host-plant  
79 blends specific to monitor *G. molesta*, several field and laboratory studies show that host-  
80 plant-released volatiles elicit male and female *G. molesta* responses that are comparable  
81 to the natural host. In Australia, a synthetic volatile blend that mimics those emitted by  
82 young peach shoots captured up to 130 males per trap, although it did not capture  
83 females (Il'ichev et al. 2009). In China, several synthetic volatile blends mimicking pear  
84 (*Pyrus bretschneideri* Rehder and *Pyrus pyrifolia* (Burm.), and peach (*Prunus persica* (L.))  
85 fruit and shoots have been identified (Lu et al. 2012, 2014, 2015). The volatile blend

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 86 obtained from pear (*P. bretschneideri* var. Jimi) fruit captured about 50 males and 20  
12  
13 87 females per trap. In the case of males, this was only 5 times less than what commercial sex  
14  
15 88 pheromone traps captured in that study (Lu et al. 2012). In addition, this blend resulted in  
16  
17 89 approximately 80% approach and 10% source contact by males in the flight tunnel; which  
18  
19 90 was equivalent to the response of the natural fruit (Lu et al. 2012). In another study in  
20  
21 91 Switzerland, analysis of peach shoot volatiles resulted in a blend that in dual-choice  
22  
23 92 olfactometer tests, was as attractive to mated females as the natural host-plant blend  
24  
25 93 (Piñero and Dorn 2007).

26  
27 94 Captures of *G. molesta* in the Australian and Chinese studies was remarkable and  
28  
29 95 paralleled the attraction of *C. pomonella* to the pear ester (Light et al. 2001). Thus, these  
30  
31 96 new blends could be an invaluable tool for the management of *G. molesta*. However, *G.*  
32  
33 97 *molesta* is a widely distributed species with significant genetic differentiation among  
34  
35 98 world populations (Kirk et al. 2013), and therefore it is crucial to determine if the  
36  
37 99 Australian and Chinese blends, which have been tested only in these two countries, are  
38  
39 100 also attractants in other areas of the geographical distribution of this species. In addition,  
40  
41 101 it would be useful to determine if the two blends are equally attractive. The Swiss blend,  
42  
43 102 which attracts females under laboratory conditions, remains to be tested in the field. The  
44  
45 103 first objective of our study was to compare the attractiveness of the Australian, Chilean  
46  
47 104 and Swiss blends and to test them at a new location. The three plant blends, plus a new  
48  
49 105 "Total" blend made with all the different components from the other three test blends,  
50  
51 106 were tested together in a peach orchard in Chile, a country where *G. molesta* was first  
52  
53 107 recorded in 1970 (González 2003). The second objective of this study was to explore the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 108 potential of the host-plant blends to increase captures of males in sex pheromone traps,  
12  
13 109 as this could improve the monitoring of males under mating disruption (Yu et al. 2014). To  
14  
15 110 this end, we compared captures in traps baited with the pheromone alone and traps  
16  
17 111 baited with the pheromone combined with the host-plant volatiles. Previous flight-tunnel  
18  
19 112 studies show that the Chinese blend attracts males and females (Lu et al. 2012), and the  
20  
21 113 Swiss blend does not attract males but synergizes their response to the sex pheromone  
22  
23 114 (Varela et al. 2011). Our third objective was to compare responses between field and  
24  
25 115 laboratory settings. To do so, the plant blends were tested in the flight tunnel alone and in  
26  
27 116 combination with the sex pheromone.

28  
29 117

## 30 31 118 Materials and Methods

### 32 33 119 *Chemical stimuli*

34  
35 120 Host-plant volatiles were purchased from Sigma-Aldrich (Santiago, Chile, chemical purity,  
36  
37 121 product and lot numbers in Table 1). The composition of the host-plant blends followed  
38  
39 122 those reported by the Australian (Il'ichev et al. 2009), Chinese (Lu et al. 2012), and Swiss  
40  
41 123 (Piñero and Dorn 2007) studies (Table 2). The Australian study used (*E*)- $\beta$ -farnesene and  
42  
43 124 (*E*)- $\beta$ -ocimene, but pure isomers were not available to us at the time of the study, so a  
44  
45 125 mixture of farnesene isomers and  $\beta$ -ocimene isomers were used instead (Table 1). A  
46  
47 126 fourth host-plant blend made by combining the components from all three blends was  
48  
49 127 included in the tests ("Total" blend, Table 2). To duplicate Australian and Chinese studies,  
50  
51 128 all chemicals were dissolved in *n*-hexane and loaded in rubber septa (red, i.d x o.d., 3.4  
52  
53 129 mm x 6.6 mm, Sigma-Aldrich, Santiago, Chile, product number Z565709), at approximately

similar concentrations as in the original studies (100 mg total and 100 mg major compound, respectively, Table 3). The Swiss blend was prepared the same way as the Chinese blend (100 mg major compound), and the Total blend was made with 10 mg of each compound. Plant odor loads for all four blends ranged between 100 and 189 mg (Table 3). With the aim of providing a wider range of release rates, in experiment 1 (see below) the host-plant blends were prepared at an additional 10-fold lower concentration, and both high and low concentrations were further dissolved in mineral oil (Sigma-Aldrich product number M8410, lot number MKBG7544V, CAS number 8020-83-5) and loaded in 1.5 mL microcentrifuge tubes fitted with a 15-mm-long x 7-mm-diameter section of dental cotton wick to absorb the chemicals. Mineral or paraffin oil is a mixture of *n*-alkanes and provide slower and more linear release rates than individual shorter alkanes, like *n*-hexane, and both are used routinely to deliver plant volatiles in olfactory tests (Andersson et al. 2012). The lids of the microcentrifuge tubes were perforated with a 1.5-mm diameter hole and were kept closed. Rubber septa were rinsed in *n*-hexane and then in acetone, and allowed to dry, before use. Microcentrifuge tubes were rinsed in acetone. Host-plant volatiles were loaded in 500  $\mu$ l volumes in rubber septa and microcentrifuge tubes at the concentrations indicated below. It took about 1 h for the host-plant blends to be absorbed by the rubber septa. For 2 to 4 h the dispensers were maintained in a well ventilated area and then placed inside plastic bottles and stored at -20C° until taken to the field on the following day.

In the Chinese study (Lu et al. 2012) commercial sex pheromone rubber septa were used. In our study the three sex pheromone components, (*Z*)-8-dodecenyl acetate (*Z*8-



12:Ac), E8-12:Ac, and (Z)-8-dodecenol (Z8-12:OH) (Pherobank, Wageningen, The Netherlands, > 99% pure) were diluted in *n*-hexane at a 100:5.4:10 ratio, respectively (Knight et al. 2015), and loaded in red rubber septa. In the first experiment we used 80 µg, which is an optimal quantity (Knight et al. 2015). Captures were relatively high so in experiments 2 and 3 we reduced the quantity of pheromone to 16 and 8 µg, respectively. This reduction in pheromone load permitted to test potential synergistic effects of host-plant volatiles on sex pheromone attraction.

#### *Field tests*

Experiments were carried out in a peach (*Prunus persica* var. *persica* cv. Doctor Davis and Carson) orchard in Chile (Duao, Maule, 35°33'29"S, 71°33'44"W) between December 21<sup>st</sup> 2012 and February 25<sup>th</sup> 2013 (Table 3). The Chinese and Australian studies used standard delta traps and funnel-type Efekto fly traps, respectively (Il'ichev et al. 2009, Lu et al. 2012). We used white delta traps (215 mm long x 200 mm wide x 100 mm tall, 340 cm<sup>2</sup> adhesive base area, Plastic Delta Trap, Alphascent, West Linn, OR, USA), except for the pheromone treatment in experiment 1, where the traps were red due to a temporary shortage of white traps. This should not have influenced trap catches because trap color does not affect *G. molesta* captures (Zhao et al. 2013). Traps were placed at 1.7 m high, hanging from 4-cm-diameter blue PVC pipes fitted in the tree branches. Traps within a plot were placed in a transect 15 to 20 m apart, and plots were at least 15 m apart from each other. Trap floors were lined with removable sticky cards.

In experiment 1 the dispensers, either rubber septa or microcentrifuge tubes, were hung from the ceiling of the trap with a wire, almost touching the trap floor. In the other two experiments the dispensers were placed directly on the sticky floor. Septa and microcentrifuge tubes were labeled with the treatment name using permanent markers. Traps lured with sex pheromone and host-plant odors (experiments 2 and 3) had 2 septa, one for each stimulus type, which were placed within a few cm of each other at the center of the trap. Sticky bottoms were replaced if there were captures. Sex of captured individuals was determined in the laboratory using a stereo microscope.

Experiment 1 was carried out between December 21<sup>st</sup> 2012 and January 29<sup>th</sup> 2013 and tested the 4 host-plant blends at two doses with two solvents and dispensers (Table 3). Sex pheromone (80  $\mu$ g) and solvent (*n*-hexane or mineral oil) were the positive and negative controls, respectively, and were loaded in the corresponding dispensers (septum or microcentrifuge tubes). The 20 plant-volatile treatments [(4 plant treatments x 2 doses + solvent + pheromone) x 2 dispenser types (septum or microcentrifuge tube)] were placed in each of 4 rows, or plots, at random. There were 6 weekly trap checks. Sex pheromone and host-plant lures were replaced on the first and second checks. For the remainder of the experiment the pheromone lure was unchanged, while new plant lures were replaced one last time on the 4<sup>th</sup> check (January 11<sup>th</sup> 2013).

Experiment 2 was ran between January 29<sup>th</sup> and February 12<sup>th</sup> 2013 to determine if the host-plant blends had any effect on male attraction to the sex pheromone. Traps were baited with a sex pheromone septum (16  $\mu$ g) and a host-plant blend septum at a similar dose as in the original Australian and Chinese studies, which was the same as the "high"

dose of experiment 1. This dose resulted in pheromone: host-plant volatile ratios ranging between 1:1,650 (Australian blend) and 1:11,812 (Chinese blend) (Table 3). Sex pheromone-only traps served as positive controls and *n*-hexane-only traps served as negative controls. The 10 treatments (4 host-plant blends; 4 sex pheromone + host-plant blends; solvent; sex pheromone) were replicated in 8 plots at random. There were 5 trap checks every 3 to 4 days. Sex pheromone septa were not replaced, and new plant lures were replaced on February 5<sup>th</sup> 2013.

Experiment 3, which was carried out between February 12<sup>th</sup> and 25<sup>th</sup> 2013, tested if the inhibitory effect of the host-plant blends observed in experiment 2 (see Results) was dose-dependent. It included the two plant blends that caused the strongest inhibition in experiment 2 (Chinese and Total, see Results), and sex pheromone at 8  $\mu$ g. The pheromone:plant ratio ranged from the lowest 1:175 (Total) to the highest 1:23,625 (Chinese) (Table 3). Two host-plant volatiles, terpinyl acetate and  $\beta$ -ocimene isomer mix, which have shown behavioral activity in previous studies (Cichón et al. 2013; Il'ichev et al. 2009; Knight et al. 2014), were tested alone at 3 mg, as in Knight et al. (2014), and in combination with sex pheromone at a sex pheromone: host-plant volatile ratio of 1:375 (Table 3). In addition, we tested the effect of having one or two septa in the trap, by adding a blank septum to a trap with a pheromone septum. *n*-Hexane septa were tested alone to control for possible sex pheromone contamination. The 13 treatments (2 host-plant blends with sex pheromone x 3 doses; sex pheromone alone; sex pheromone and *n*-hexane septum; *n*-hexane; 2 host-plant volatiles with and without sex pheromone) were placed in 4 plots at random. There were 6 checks every 1 or 2 days because there was a

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 217 population peak during this period. Pheromone-loaded septa were not replaced, yet new  
12  
13 218 host-plant lures were replaced on February 18<sup>th</sup> and 22<sup>nd</sup> 2013.  
14  
15 219  
16  
17 220 *Flight tunnel test*  
18  
19 221 The colony of *G. molesta* used in the flight tunnel was established with insects collected at  
20  
21 222 Piacenza, Italy. The population has been maintained at the University of Lleida, Spain,  
22  
23 223 since 2005 without reintroduction of wild individuals. Larvae were reared on a semi-  
24  
25 224 synthetic diet modified from Ivaldi-Sender (1974) under a L16:D8 photoperiod at  $24 \pm 1^\circ\text{C}$ .  
26  
27 225 Pupae were separated by sex and placed in 4-L polypropylene containers and provided a  
28  
29 226 cotton ball soaked in 10% sugar water solution. Adults were collected daily and used when  
30  
31 227 2 to 4 days old.  
32  
33 228 The flight tunnel and its methodology have been previously described  
34  
35 229 (Ammagarahalli et al. 2017). The 150 x 45 x 45 cm (length x height x width) tunnel had a  
36  
37 230  $0.35 \text{ m s}^{-1}$  wind flow and the temperature was maintained at  $23 \pm 1^\circ\text{C}$ . It was illuminated  
38  
39 231 from above with 36-watt fluorescent lamps producing 150 lux white light. Tests were  
40  
41 232 carried out during the last 3 hours of the photophase and occasionally into the first hour  
42  
43 233 of the scotophase, in which case the daylight illumination was left on. Males were placed  
44  
45 234 individually in glass tubes and were transferred to the flight tunnel room 30 to 120 min  
46  
47 235 before the beginning of the test. Test odors were applied in 10  $\mu\text{l}$  loads to 10 x 15 mm  
48  
49 236 filter paper pieces that were allowed to dry for 5 to 10 min until tested in the flight tunnel  
50  
51 237 5 to 180 min later. The male was placed in the flight tunnel after the odor stimulus, on top  
52  
53 238 of a metal-wire platform similar to the one used for the odor source and 1.3 m downwind  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 239 from it. We recorded for 2 min if the male took flight, started upwind oriented flight  
12  
13 240 (zigzagging upwind flight) or landed on the filter paper containing the stimulus source, and  
14  
15 241 the time it took the males to engage in these behaviors. The stimulus was placed in the  
16  
17 242 tunnel on top of a 25-cm-tall metal-wire platform. Three to five males flew to each filter  
18  
19 243 paper treatment before changing to another treatment paper. At the end of a test day a  
20  
21 244 filter paper had been used with 8 to 15 males, therefore, the filter papers were outside  
22  
23 245 their individual glass vial and exposed to the wind flow between of 32 to 60 min before  
24  
25 246 being discarded. On any given day, only one filter paper was used for each treatment. Due  
26  
27 247 to the high number (see below), the treatment order was randomized in two groups and  
28  
29 248 tested on alternate days.

30  
31 249 The following treatments were tested in the flight tunnel: Australian, Chinese and  
32  
33 250 Swiss host-plant blends at 10  $\mu$ g each, a suboptimal sex pheromone dose of 1 ng (a  
34  
35 251 response curve to doses of 1, 10, 100 and 1,000 ng resulted in 34, 82, 89 and 63% source  
36  
37 252 contact respectively, N = 44, similar to Ammagarahalli et al. (2017)), and sex pheromone:  
38  
39 253 host-plant blends (Australian, Chinese or Swiss) at 1:10, 1:100, 1:1,000, and 1:10,000  
40  
41 254 ratios with sex pheromone at 1 ng (Table 3). We used a suboptimal dose of sex  
42  
43 255 pheromone because the optimal dose results in a high percentage of response. The 10  $\mu$ g  
44  
45 256 dose of host-plant odors was 50 times lower than that used in the Chinese flight tunnel  
46  
47 257 test with rubber septa (Lu et al. 2012), yet similar to what we have used previously (Varela  
48  
49 258 et al. 2011). To control contamination, 20 insects were tested with *n*-hexane on random  
50  
51 259 days. The sex pheromone:host-plant blends were prepared using a stock sex pheromone  
52  
53 260 solution so all had identical sex pheromone concentrations. The 1:0, 1:100, 1:1,000 and

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 261 1:10,000 blends were prepared on January 18<sup>th</sup>, 2013 and the 1:10 blend on February 4<sup>th</sup>,  
12  
13 262 2013. Flight tunnel tests were carried out between February 8<sup>th</sup> and 27<sup>th</sup>, 2013 with N =  
14  
15 263 64.  
16  
17 264  
18  
19 265 *Statistical analyses*  
20  
21 266 Generalized linear models (GLM) with Poisson family function in the package lme4 of R (R  
22  
23 267 Development Core Team 2015) were used to analyze trap count data (Bolker et al. 2009).  
24  
25 268 Due to the high temporal variation in trap captures, sampling date was included as a  
26  
27 269 random effect in the generalized linear mixed model (GLMM), along with the variation  
28  
29 270 among plots if they contributed significantly to the model after comparing among models  
30  
31 271 with ANOVA. To treatments with zero captures, a random capture was added so the  
32  
33 272 GLMM could converge. The percentage of males which responded in the flight tunnel was  
34  
35 273 analyzed with GLM models using a binomial family function. Behavioral categories (take  
36  
37 274 flight, oriented flight and contact) were analyzed separately. One response was added to  
38  
39 275 treatments with no responses in a randomly chosen replicate so the GLM model could  
40  
41 276 converge. The time elapsed before insect response inside the flight tunnel was analyzed  
42  
43 277 with a linear model, lm(), in transformed [log (x+1)] data. Comparisons among treatment  
44  
45 278 pairs in both field and flight tunnel studies were performed with the glht() or lsmeans()  
46  
47 279 functions of R using Tukey's alpha correction method. Whenever the term "significant" is  
48  
49 280 used in the text regarding treatment comparisons it indicates that  $p \leq 0.05$ . Raw data and  
50  
51 281 R scripts are available at <https://repositori.udl.cat/handle/10459.1/59534>  
52  
53 282  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10 283 Results  
11  
12  
13 284 *Field tests*  
14  
15 285 In experiment 1 only one female was captured in a trap baited with a high dose of the  
16  
17 286 Swiss host-plant blend diluted in mineral oil. The four traps baited with sex pheromone  
18  
19 287 septa captured a total of 1,632 males, whereas, the four traps baited with sex pheromone  
20  
21 288 microcentrifuge tubes captured a total of 215 males. All of the host-plant volatile traps  
22  
23 289 combined, which summed 64, captured a total of 64 males in the entire experiment (Table  
24  
25 290 4). Of these 64 males, 61 were captured in the 4<sup>th</sup> weekly check, and these captures  
26  
27 291 clustered mainly in 3 particular traps: 35 males in a trap baited with the low-dose Swiss  
28  
29 292 host-plant blend in *n*-hexane, 15 males in a trap baited with the high-dose Chinese host-  
30  
31 293 plant blend in mineral oil, and 9 males in a trap baited with a high-dose Total host-plant  
32  
33 294 blend in *n*-hexane. This level of captures in host-plant baited traps was not observed  
34  
35 295 before or after week 4 (only 3 more males were captured by host-plant-baited traps in the  
36  
37 296 other 5 weekly checks) or in experiment 2 (see below). Table 4 summarizes total trap  
38  
39 297 captures, and shows how the Swiss host-plant blend at the low dose in *n*-hexane was the  
40  
41 298 only host-plant blend that captured significantly more males ( $1.46 \pm 1.46$   
42  
43 299 males/trap/check, mean  $\pm$  SEM), than the *n*-hexane traps, which captured none.  
44  
45 300 In experiment 2, traps baited with host-plant volatiles or with *n*-hexane captured  
46  
47 301 no males, whereas sex pheromone-baited traps captured many males (2,800 in total,  
48  
49 302 Table 5). Only 9 females were captured in this experiment, all of them in traps baited with  
50  
51 303 the Australian host-plant blend combined with the sex pheromone, but these captures  
52  
53 304 were not significantly different than those from the *n*-hexane traps. The addition of a  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 305 septum baited with any host-plant blend to a trap baited with a sex pheromone septum  
12  
13 306 significantly decreased the number of males captured with respect to sex pheromone  
14  
15 307 traps. This negative effect was significantly stronger for the Chinese and Total host-plant  
16  
17 308 blends than for the Australian and Swiss host-plant blends.

18  
19 309 In experiment 3, a total of 13,650 males and 17 females were captured. The  
20  
21 310 addition of a septum baited with any of the three doses of the Chinese or Total host-plant  
22  
23 311 blends to a trap baited with a sex pheromone septum significantly decreased the number  
24  
25 312 of males captured relative to traps baited with a sex pheromone septum and an *n*-hexane  
26  
27 313 septum. This effect was more pronounced as the host-plant dose increased (Table 6).  
28  
29 314 Traps baited with a sex pheromone septum and an *n*-hexane septum captured more  
30  
31 315 males than traps baited with only the sex pheromone septum. *n*-Hexane traps captured 6  
32  
33 316 males in total.

34  
35 317  $\beta$ -ocimene (mixture of isomers) captured significantly more males than *n*-hexane  
36  
37 318 traps. Terpinyl acetate added to sex pheromone significantly increased captures relative  
38  
39 319 to traps baited with a sex pheromone septum, but not relative to traps baited with a sex  
40  
41 320 pheromone and a *n*-hexane septum together (Table 6).

42  
43 321 The 17 females caught in experiment 3 were captured by 8 different treatments, 5  
44  
45 322 of which captured only one female, while 3 captured more than one female. The  
46  
47 323 treatments that captured more than 1 female were always a combination of the host-  
48  
49 324 plant blend with the sex pheromone. Of these, the highest captures were in  $\beta$ -ocimene  
50  
51 325 (mix of isomers) plus sex pheromone which captured 3 females in one plot on three  
52  
53 326 different dates, 3 females in another plot on two different dates, and one female in



another plot. None of the female captures in these treatments were significantly higher than *n*-hexane traps.

#### *Flight tunnel test*

In the flight tunnel experiment several doses of the Australian, Chinese and Swiss host-plant blends were added to sex pheromone. None of the host-plant blends alone, nor *n*-hexane, attracted any males, so they were not included in the means comparison test. Pairwise comparisons between each sex pheromone host-plant combination treatment and the isolated sex pheromone treatment showed that the three host-plant blends significantly increased the percentages of flight, oriented flight and contact, and did so in a dose-dependent manner (Fig. 1a). The Chinese and Swiss host-plant blends significantly increased responses at the 1:100 to 1:1,000 sex pheromone:host-plant ratios, whereas the Australian host-plant blend did so at the 1:1 and 1:10 ratios. The three host-plant blends significantly reduced the time of response to the sex pheromone, and, as with the percentages of response, the effect was stronger at the higher (plant wise) sex pheromone:host-plant odor ratios (1:1,000 and 1:10,000) (Fig. 1b).

#### Discussion

Captures by the Australian and Chinese host-plant blends in our study were substantially lower than in the original studies (from here onwards "Australian and Chinese studies" will refer to "Il'ichev et al. 2009 and Lu et al. 2012", respectively). Although the experimental conditions of the Australian and Chinese studies were closely reproduced in Chile, there

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

349 were some differences that perhaps account for the low response of these host-plant  
350 blends in our experiments. The most evident difference is the chemical purity of farnesene  
351 and ocimene, which in the Australian study consisted of (*E*)- $\beta$ -farnesene and (*E*)- $\beta$ -  
352 ocimene. The isolated isomers were not commercially available at the time of the tests,  
353 therefore, a mixture of isomers of each compound were used in our study. Insects can  
354 distinguish odorant isomers both at the sensory and behavioral levels (De Bruyne and  
355 Baker 2008), so it is possible that isomeric purity affected reproducibility of the Australian  
356 blend in Chile. However, the  $\beta$ -ocimene mix of isomers in our study was one of the few  
357 plant stimuli that was more attractive than *n*-hexane, whereas the Australian blend  
358 influenced male response to pheromone, as did the other plant blends. Although our  
359 Australian blend did not have the same isomer purity as the original Australian blend, our  
360 results show that it was sensed by and affected the behavior of *G. molesta*. Furthermore,  
361 the Chinese blend used a mixture of farnesene isomers both in the original Chinese study  
362 and in our study. Even though it was very attractive in China, it performed very poorly in  
363 Chile. Therefore, isomer purity should not have contributed to the different performance  
364 of the Chinese blend in China and Chile. Little is known about the detection of plant  
365 volatiles in *G. molesta*, but olfactory receptor neurons relatively specific to the same  
366 farnesene isomer mix that we used in here have been described on the antenna of males  
367 (Ammagarahalli and Gemeno 2015).

368         Although we used the same solvents, concentrations and dispensers as in the  
369 original Australian and Chinese studies, the quantity and proportion of odorants released  
370 by the dispensers were not analyzed in any of these studies. Due to this we do not know if

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

the volatile composition and emission rate would have varied among them. By using an additional lower concentration than in the original Australian and Chinese studies, and mineral oil as an extra solvent, we attempted to diversify the stimulus quantity released by the dispensers. Mineral oil probably decreased release rates because pheromone in mineral oil attracted about 7-times fewer males than pheromone in *n*-hexane. A similar solvent effect was probably true for the host-plant volatiles, however neither mineral oil nor the lower stimulus concentration improved captures relative to the original solvent and concentration. In our tests, dispensers with plant volatiles were loaded a day before used and replaced every  $5.6 \pm 3.9$  days (mean  $\pm$  SEM) in order to minimize stimulus degradation and depletion during the assay. A final methodological difference among studies was the use of a funnel-type fly trap in the Australian study, whereas in the Chinese and Chilean studies standard delta traps were used. Although trap type could affect moth captures, our sex pheromone traps captured many males, thus it is unlikely that trap type alone could explain the dissimilar performance of the Australian blend in Australia and Chile. With regard to the flight tunnel test, we used a lower host-plant odor concentration than the Chinese study, and loaded it on filter paper instead of rubber septa, so the difference between the lack of response in our flight tunnel test and the good response in the Chinese flight tunnel test could be related to the use of different stimulus delivery conditions.

Biological variables such as population genetics could also be involved in the differences between the Australian and Chinese studies and our Chilean test. When populations evolve host-specialization, differences in the detection of host-plant volatile

signals may arise (Smadja and Butlin 2009). Significant genetic differences among populations of *G. molesta* could provide the necessary genetic diversity for the evolution of host varieties (Zheng et al. 2015; Wei et al. 2015). Nonetheless, host-plant specialization may be negligible in *G. molesta* given its recent human-aided expansion and reduced dispersal power (Wei et al. 2015). This is consistent with the lack of worldwide variation in sex pheromone production and response that we have reported in this species (Knight et al. 2015). However, whether this is also true regarding responses to host-plant volatiles remains to be tested.

Background odors in the environment could influence the response of insects to odor stimuli (Cai et al. 2017). Pear ester is more effective in attracting *C. pomonella* when deployed in walnut orchards than in pear orchards (Light et al. 2001). In China, pear- and peach-odor blends perform relatively better in orchards of the opposite host (Lu et al., 2014; 2015). Captures with a pear blend in a peach orchard were >40 males per trap (Fig 2A in Lu et al. 2015). With this in mind we expected substantial captures of *G. molesta* in our peach fields with the Chinese pear blend, but captures were never higher than in control traps. The Australian peach blend was very attractive in a pear background (Il'ichev et al., 2009), so its poor performance in our peach orchard might be related to the differences in host-plant backgrounds. Yet, in China the peach blends attract *G. molesta* in peach orchards (>10 males/trap, Fig 2A in Lu et al. 2015), and the pear blend is equally effective in peach and pear orchards (Fig 2B in Lu et al., 2012), so the relative importance of background odors needs to be examined. Background odors, however, could be relatively important to explain differences between laboratory and field studies. Knudsen

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 415 et al. (2008), for example, show the host-plant volatile compound which attracts the apple  
12  
13 416 fruit moth, *Argyresthia conjugella* Zeller, in the flight tunnel is different than the one that  
14  
15 417 attracts it in the field, albeit these compounds are released by the host. The authors  
16  
17 418 conclude that the interaction of the plant volatiles with the background odor contributes  
18  
19 419 to different results between field and laboratory tests. The flower volatile  
20  
21 420 phenylacetaldehyde is a generalist noctuid moth attractant (Tóth et al. 2010) that  
22  
23 421 increases the response of male *Spodoptera frugiperda* Walker to sex pheromone in the  
24  
25 422 flight tunnel (Meagher and Mitchel 1998). However, this flower volatile decreases  
26  
27 423 captures in sex pheromone traps in the field (Meagher 2001). Perhaps the lack of  
28  
29 424 background plant odors in our flight tunnel test could explain why males were more  
30  
31 425 attracted to the pheromone plus host-plant lures than to the pheromone lure alone.  
32  
33 426 However, it seems unlikely that the presence of background plant odors in the field is  
34  
35 427 responsible for the inhibitory effect of the host-plant blends on sex pheromone traps.  
36  
37 428 An important advantage of host-plant volatiles over sex pheromones is the  
38  
39 429 potential attraction of females, which provides information on the mating status of the  
40  
41 430 population. The Australian study reports that only males were attracted to the plant lures  
42  
43 431 (Il'ichev et al. 2009), and the Chinese study reports male:female ratios in the range of 10:1  
44  
45 432 to 3:1 (Lu et al. 2012). In our study the plant odor blends also caught substantially more  
46  
47 433 males than females (122 vs 2 in total, respectively). Interestingly, 24 additional females  
48  
49 434 (92% of all the females caught in the study) were collected in traps baited with a  
50  
51 435 combination of sex-pheromone and host-plant septa. Given recent reports of pheromone  
52  
53 436 autodetection by female moths, and its possible implications in mating disruption

(Holdcraft et al. 2016), it may be worth exploring if the sex pheromone plays any role on female moth attraction to host-plant lures.

A field study with *G. molesta* shows that two plant volatiles [(Z)-3-hexenyl acetate and undecanol] added to the sex pheromone synergize captures at a 1:0.5 pheromone:host-plant ratio, but at the 1:1 and 1:2 ratios the synergism disappears, with a clear trend to become inhibitory at even higher host-plant ratios (Yu et al. 2014). In other moth species where synergism of host-plant volatiles with sex pheromone has been reported, the pheromone:host-plant ratio tested is in the range of 1:1 (e.g., Dai et al. 2008; Dickens et al. 1993; Light et al. 1993), although sometimes it can be as high as 1:40 (Deng et al. 2004). We used much higher sex pheromone:host-plant ratios, around 1:10,000, in order to maintain the same host-plant concentrations as in the original Australian and Chinese studies. But when we later tested a lower (1:200) ratio in the dose-response test (experiment 3), the effect was also inhibitory relative to traps baited with a sex pheromone and an *n*-hexane septum. Thus, it remains to be determined if still lower sex pheromone:host-plant ratios of the Australian and Chinese plant-blends could have a non-inhibitory, or even a synergistic effect on sex pheromone captures. There are at least two other cases where host-plant odors reduce moth captures in sex pheromone traps. (E)-2-Penten-1-ol reduced captures of the arctiid moth *Hyphantria cunea* (Drury) when added to the sex pheromone at a 1:1 ratio (Tang et al. 2012), and phenylacetaldehyde reduced pheromone captures of *Spodoptera frugiperda* (Smith) in traps baited with a commercial sex pheromone septum and different amounts of the host-plant odorant (Meagher 2001).

Increased captures by the presence of a blank septum in sex pheromone traps was unexpected, and could be a visual response because *G. molesta* flies during the last light hours of the day using visual information (Kuenen and Gilbert 2014). It could also be a response to increased air turbulence, a factor that facilitates male upwind flight (De Bruyne and Baker 2008). Barros-Parada et al. (2016) report a large effect of the location of pheromone and pear ester septa and acetic acid dispensers in the trap on captures of *C. pomonella*. Our observations highlight the importance of taking into account apparently negligible experimental variables, like the addition of a second septum, which could impact trap captures and hamper the correct evaluation of host-plant attractants.

#### Acknowledgments

B.A. acknowledges support by a Ph.D. fellowship from MINECO (Spain). W.B.P acknowledges a FONDECYT-Postdoctorado grant 3140285 (Chile). The visit of C.G. to Chile was funded by Chilean Government grant CONICYT-Atracción de Capital Humano Avanzado del Extranjero - Modalidad Estadías Cortas (MEC) 80120005. Field work in Chile was supported by Alexis Muñoz, Nicolás Berríos and Carlos Cavieres.

#### References

Allison JD, Cardé RT (2016) Pheromones: Reproductive isolation and evolution in moths. In: Allison JD and Cardé RT (eds) Pheromone communication in moths: Evolution, behavior and application. University of California Press, Oakland, pp 11–24

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 480 Ammagarahalli B, Gemenó C (2015) Interference of plant volatiles on pheromone receptor  
12  
13 481 neurons of male *Grapholita molesta* (Lepidoptera: Tortricidae). J Insect Physiol  
14  
15 482 81:118–128. doi: 10.1016/j.jinsphys.2015.07.009  
16  
17 483 Ammagarahalli C, Chianella L, Gomes P, Gemenó C (2017) Role of plant volatiles and  
18  
19 484 hetero-specific pheromone components in the wind tunnel response of male  
20  
21 485 *Grapholita molesta* (Lepidoptera:Tortricidae) to modified sex pheromone blends.  
22  
23 486 Bull Entomol Res 107:573-582. doi: 10.1017/S0007485317000013  
24  
25 487 Andersson MN, Schlyter F, Hill SR, Dekker T (2012) What reaches the antenna? How to  
26  
27 488 calibrate odor flux and ligand–receptor affinities. Chem Senses 37:403–420. doi:  
28  
29 489 10.1093/chemse/bjs009  
30  
31 490 Barros-Parada W, Basoalto E, Fuentes-Contreras E, Cichón L, Knight AL (2016) Acetic acid  
32  
33 491 lure placement within traps affects moth catches of codling moth (Lepidoptera:  
34  
35 492 Tortricidae). J Appl Entomol 140:786–795. doi: 10.1111/jen.12311  
36  
37 493 Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White JSS (2009)  
38  
39 494 Generalized linear mixed models: a practical guide for ecology and evolution.  
40  
41 495 Trends Ecol Evol 24:127–135. doi: 10.1016/j.tree.2008.10.008  
42  
43 496 Cai X, Bian L, Xu X, Luo Z, Li Z, Chen Z (2017) Field background odour should be taken into  
44  
45 497 account when formulating a pest attractant based on plant volatiles. Sci Rep  
46  
47 498 7:41818. doi: 10.1038/srep41818  
48  
49 499 Cichón L, Fuentes-Contreras E, Garrido S, Lago J, Barros-Parada W, Basoalto E, Hilton R,  
50  
51 500 Knight AL (2013) Monitoring oriental fruit moth (Lepidoptera: Tortricidae) with



501 sticky traps baited with terpinyl acetate and sex pheromone. J Appl Entomol  
 502 137:275-281. doi: 10.1111/j.1439-0418.2012.01732.x  
 503 Dai J, Deng J, Du J (2008) Development of bisexual attractants for diamondback moth  
 504 *Plutella xylostella* (Lepidoptera: Plutellidae) based on sex pheromone and host  
 505 volatiles. Appl Entomol Zool 43:631–638. doi: 10.1303/aez.2008.631  
 506 De Bruyne M, Baker TC (2008) Odor detection in insects: volatile codes. J Chem Ecol  
 507 34:882–897. doi: 10.1007/s10886-008-9485-4  
 508 Deng JY, Wei HY, Huang YP, Du JW (2004) Enhancement of attraction to sex pheromones  
 509 of *Spodoptera exigua* by volatile compounds produced by host plants. J Chem Ecol  
 510 30:2037–2045. doi: 10.1023/B:JOEC.0000045593.62422.73  
 511 Dickens JC, Smith JW, Light DM (1993) Green leaf volatiles enhance sex attractant  
 512 pheromone of the tobacco budworm *Heliothis virescens* (Lep: Noctuidae).  
 513 Chemoecology 4:175–177. doi: 10.1007/BF01256553  
 514 González RH (2003). Las polillas de la fruta en Chile (Lepidoptera: Tortricidae; Pyralidae).  
 515 Universidad de Chile. [in Spanish with English summary]  
 516 Guillelme LJ, Iguchi T (2012) Life in a contaminated world. Science 337:1614–1615. doi:  
 517 10.1126/science.1226985  
 518 Holdcraft R, Rodriguez-Saona C, Stelinski LL (2016) Pheromone autodetection: evidence  
 519 and implications. Insects 7:17. doi: 10.3390/insects7020017  
 520 Il'ichev AL, Kugimiya S, Williams DG, Takabayashi J (2009) Volatile compounds from young  
 521 peach shoots attract males of oriental fruit moth in the field. J Plant Interact  
 522 4:289–294. doi: 10.1080/17429140903267814

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11 523 Ivaldi-Sender C (1974) Techniques simples pour un élevage permanent de la tordeuse
- 12
- 13 524 orientale *Grapholita molesta* (Lepidoptera: Tortricidae) sur milieu artificiel. Ann
- 14
- 15 525 Zool Ecol Anim 6:337–343. [in French with English summary]
- 16
- 17 526 Kirk H, Dorn S, Mazzi D (2013) Worldwide population genetic structure of the oriental fruit
- 18
- 19 527 moth (*Grapholita molesta*), a globally invasive pest. BMC Ecology 13:1–12. doi:
- 20
- 21 528 10.1186/1472-6785-13-12
- 22
- 23 529 Knight A, Basoalto E, Hilton R, Molinari F, Zoller B, Hansen R, Crawczyk, Hull L (2013)
- 24
- 25 530 Monitoring oriental fruit moth (Lepidoptera: Tortricidae) with the Ajar bait trap in
- 26
- 27 531 orchards under mating disruption. J Appl Entomol 137:650–660. doi:
- 28
- 29 532 10.1111/jen.12061
- 30
- 31 533 Knight AL, Cichón L, Lago J, Fuentes-Contreras E, Barros-Parada W, Hull L, Krawczyk G,
- 32
- 33 534 Zoller B, Hansen R, Hilton R, Basoalto E (2014) Monitoring oriental fruit moth and
- 34
- 35 535 codling moth (Lepidoptera: Tortricidae) with combinations of pheromones and
- 36
- 37 536 kairomones. J Appl Entomol 138:783–794. doi: 10.1111/jen.12138
- 38
- 39 537 Knight AL, Barros-Parada W, Bosch D, Escudero-Colomar LA, Fuentes-Contreras E,
- 40
- 41 538 Hernández-Sánchez J, Yung C, Kim Y, Kovanci OB, Levi A, Lo P, Molinari F, Valls J,
- 42
- 43 539 Gemenó C (2015) Similar worldwide patterns in the sex pheromone signal and
- 44
- 45 540 response in the oriental fruit moth, *Grapholita molesta* (Lepidoptera:Tortricidae).
- 46
- 47 541 Bull Entomol Res 105:23–31. doi: 10.1017/S0007485314000637
- 48
- 49 542 Knudsen GK, Bengtsson M, Kobro S, Jaastad G, Hofsvang T, Witzgall P (2008) Discrepancy
- 50
- 51 543 in laboratory and field attraction of apple fruit moth *Argyresthia conjugella* to host
- 52
- 53 544 plant volatiles. Physiol Entomol 33:1–6. doi: 10.1111/j.1365-3032.2007.00592.x
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

545 Kong WN, Li J, Fan RJ, Li SC, Ma R (2014) Sex-pheromone-mediated mating disruption  
546 technology for the oriental fruit moth *Grapholita molesta* (Busck) (Lepidoptera:  
547 Tortricidae): overview and prospects. *Psyche* 2014:1–8. doi: 10.1155/2014/253924  
548 Kuenen L, Gilbert C (2014) Visual ground pattern modulates flight speed of male Oriental  
549 fruit moths. *Physiol Entomol* 39:271–279. 10.1111/phen.12072  
550 Light DM, Flath RA, Buttery RG, Zalom FG, Rice RE, Dickens JC, Jang EB (1993) Host-plant  
551 green-leaf volatiles synergize the synthetic sex pheromones of the corn earworm  
552 and codling moth (Lepidoptera). *Chemoecology* 4:145–152. doi:  
553 0.1007/BF01256549  
554 Light DM, Knight AL, Henrick CA, Rajapaska D, Lingren B, Dickens JC, Reynolds KM, Buttery  
555 RG, Merrill G, Roitman J, Bruce CC (2001) A pear-derived kairomone with  
556 pheromonal potency that attracts male and female codling moth, *Cydia pomonella*  
557 (L.). *Naturwissenschaften* 88:333–338. doi: 10.1007/s001140100243  
558 Light DM, Grant JA, Haff RP, Knight AL (2017) Addition of pear ester with sex pheromone  
559 enhances disruption of mating by female codling moth (Lepidoptera: Tortricidae)  
560 in walnut orchards treated with meso dispensers. *Environ Entomol* 46:319–327.  
561 doi: 10.1093/ee/nvw168  
562 Lu PF, Huang LQ, Wang CZ (2012) Identification and field evaluation of pear fruit volatiles  
563 attractive to the oriental fruit moth *Cydia molesta*. *J Chem Ecol* 38:1003–1016. doi:  
564 10.1007/s10886-012-0152-4

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11 565 Lu PF, Qiao HL, Xu ZC, Cheng J, Zong SX, Luo PQ (2014) Comparative analysis of peach and
- 12 pear fruit volatiles attractive to the oriental fruit moth *Cydia molesta*. J Plant
- 13 566 Interact 9:388–395. doi: 10.1080/17429145.2013.843724
- 14 567
- 15
- 16 568 Lu PF, Wang R, Wang CZ, Luo YQ, Qiao HL (2015) Sexual differences in electrophysiological
- 17 and behavioral responses of *Cydia molesta* to peach and pear volatiles. Entomol
- 18 569 Exp Appl 157:279–290. doi: 0.1111/eea.12362
- 19 570
- 20
- 21 571 Meagher Jr RL, Mitchell ER (1998) Phenylacetaldehyde enhances upwind flight of male fall
- 22 armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) to its sex pheromone.
- 23 572 Fla Entomol 81:556–559. doi: 10.2307/3495958
- 24 573
- 25 574 Meagher Jr RL (2001) Trapping fall armyworm (Lepidoptera: Noctuidae) adults in traps
- 26 baited with pheromone and a synthetic floral volatile compound. Fla Entomol
- 27 575 84:288–292. doi: 10.2307/3496181
- 28 576
- 29
- 30 577 Miller JR, Gut LJ (2015) Mating disruption for the 21<sup>st</sup> century: Matching technology with
- 31 mechanism. Environ Entomol 44:427–453. doi: 10.1093/ee/nvv052
- 32 578
- 33 579 Myers CT, Hull LA, Krawczyk G (2007) Effects of orchard host plants (apple and peach) on
- 34 development of oriental fruit moth (Lepidoptera: Tortricidae). J Econ Entomol
- 35 580 100:421–430. doi: 10.1093/jee/99.4.1176
- 36 581
- 37 582 Piñero JC, Dorn S (2007) Synergism between aromatic compounds and green leaf volatiles
- 38 derived from the host plant underlies female attraction in the oriental fruit moth.
- 39 583 Entomol Exp Appl 125:185–194. doi: 10.1111/j.1570-7458.2007.00614.x
- 40 584
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

585 R Development Core Team (2015) R: A language and environment for statistical  
 586 computing. R Foundation for Statistical Computing. Vienna Austria. ISBN 3-900051-  
 587 07-0, URL <http://www.R-project.org/>  
 588 Rothschild GHL, Vickers RA (1991) Biology, ecology and control of the oriental fruit moth.  
 589 In: van der Geest LPS, Evenhuis HH (eds) Tortricid pests: their biology, natural  
 590 enemies and control. World Crop Pests, Vol. 5. Elsevier, Amsterdam, pp 389–412  
 591 Smadja C, Butlin RK (2009) On the scent of speciation: the chemosensory system and its  
 592 role in premating isolation. *Heredity* 102:77–97. doi: 10.1038/hdy.2008.55  
 593 Szendrei Z, Rodriguez-Saona C (2010) A meta-analysis of insect pest behavioral  
 594 manipulation with plant volatiles. *Entomol Exp Appl* 134:201–210. doi:  
 595 10.1111/j.1570-7458.2009.00954.x  
 596 Tang R, Zhang JP, Zhang ZN (2012) Electrophysiological and behavioral responses of male  
 597 fall webworm moths (*Hyphantria cunea*) to herbivory-induced mulberry (*Morus*  
 598 *alba*) leaf volatiles. *PLoS one* 7:e49256. doi: 10.1371/journal.pone.0049256  
 599 Tóth M, Szarukán I, Dorogi B, Gulyás A, Nagy P, Rozgonyi Z (2010) Male and female  
 600 noctuid moths attracted to synthetic lures in Europe. *J Chem Ecol* 36:592–598. doi:  
 601 10.1007/s10886-010-9789-z  
 602 Varela N, Avilla J, Anton S, Gemenó C (2011) Synergism of pheromone and host-plant  
 603 volatile blends in the attraction of *Grapholita molesta* males. *Entomol Exp Appl*  
 604 141:114–122. doi: 10.1111/j.1570-7458.2011.01171.x  
 605 Wei SJ, Cao LJ, Gong YJ, Shi BC, Wang S, Zhang F, Guo XJ, Wang YM, Chen XX (2015)  
 606 Population genetic structure and approximate Bayesian computation analyses


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 607 reveal the southern origin and northward dispersal of the oriental fruit moth  
12  
13 608 *Grapholita molesta* (Lepidoptera: Tortricidae) in its native range. Mol Ecol  
14  
15 609 24:4094–4111. doi: 10.1111/mec.13300  
16  
17 610 Witzgall P, Kirsch P, Cork A (2010) Sex pheromones and their impact on pest management.  
18  
19 611 J Chem Ecol 36:80–100. doi: 10.1007/s10886-009-9737-y  
20  
21 612 Yu H, Feng J, Zhang Q, Xu H (2014) (Z)-3-hexenyl acetate and 1-undecanol increase male  
22  
23 613 attraction to sex pheromone trap in *Grapholita molesta* (Busck) (Lepidoptera:  
24  
25 614 Tortricidae). Int J Pest Manage 61:30–35. doi: 10.1080/09670874.2014.986560  
26  
27 615 Zhao ZG, Rong EH, Li SC, Zhang LJ, Kong WN, Hu RS, Zhang JT, Ma RY (2013) Research on  
28  
29 616 the practical parameters of sex pheromone traps for the oriental fruit moth. Pest  
30  
31 617 Manag Sci 69:1181–1186. doi: 10.1002/ps.3592  
32  
33 618 Zheng Y, Qiao X, Wang K, Dorn S, Chen M (2015) Population genetics affected by pest  
34  
35 619 management using fruit-bagging: a case study with *Grapholita molesta* in China.  
36  
37 620 Entomol Exp Appl 156:117–127. doi: 10.1111/eea.12316  
38  
39 621

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

622 **Fig. 1** Effects of adding various host-plant volatile blends (Australian, Chinese and Swiss) of  
623 different doses on the behavioral responses of *G. molesta* males to a suboptimal dose of  
624 sex pheromone in a flight tunnel (experiment 4). a) Percentage of males engaged in each  
625 behavioral category (take flight, oriented upwind flight and source contact). b) Time it  
626 took males to engage in each behavioral category. Asterisks indicate significant  
627 differences between the sex pheromone host-plant combination treatments and the  
628 isolated sex pheromone treatment (1:0) by means of planned pairwise comparisons using  
629 Tukey's test ( $p \leq 0.05$ )

Formatted: Font: Italic

Figure





**Table 1** Synthetic host-plant odorants used in the experiments

| Compound                       | CAS        | Product number (Sigma Aldrich) | Lot number | Purity <sup>a</sup> (≥ %) |
|--------------------------------|------------|--------------------------------|------------|---------------------------|
| 1-Hexanol                      | 111-27-3   | H13303                         | STBC8538V  | 98                        |
| Nonanal                        | 124-19-6   | W278203                        | STBC3506V  | 95                        |
| Ethyl butanoate                | 104-54-4   | E15701                         | STBB7416V  | 99                        |
| Butyl acetate                  | 123-86-4   | 402842                         | SHBB8826V  | 99.5                      |
| Ethyl hexanoate                | 123-66-0   | 148962                         | S28172V    | 99                        |
| Hexyl acetate                  | 142-92-7   | 108154                         | STBC6608V  | 99                        |
| Hexyl butanoate                | 2639-63-6  | W256803                        | STBC0651V  | 98                        |
| Farnesene <sup>b</sup>         | NA         | W383902                        | MKBG4494V  | NA                        |
| ( <i>Z</i> )-3-Hexenyl acetate | 3681-71-8  | W317101                        | MKBG6087V  | 98                        |
| ( <i>Z</i> )-3-Hexenol         | 928-96-1   | W256307                        | MKBG7249V  | 98                        |
| ( <i>E</i> )-2-Hexenal         | 6728-26-3  | W256005                        | STBC8608V  | 95                        |
| Benzaldehyde                   | 100-52-7   | B1334                          | STBC6885V  | 99                        |
| Benzonitrile                   | 100-47-0   | 12722                          | BCBH8265V  | 98                        |
| β-Ocimene <sup>b</sup>         | 13877-91-3 | W353901                        | MKBK5322V  | 90                        |
| Pear ester                     | 3025-30-7  | W314803                        | STBC4363V  | 80                        |
| Terpinyl acetate               | 80-26-2    |                                |            | 95                        |

<sup>a</sup> As indicated by manufacturer<sup>b</sup> Mixture of isomers

**Table 2** Odorant ratios in the four tested host-plant blends. Actual quantities used in the tests are shown in Table 3

| Plant compound                 | Host-plant blend name   |                      |                    | Total |
|--------------------------------|-------------------------|----------------------|--------------------|-------|
|                                | Australian <sup>a</sup> | Chinese <sup>b</sup> | Swiss <sup>c</sup> |       |
| 1-Hexanol                      |                         | 1                    |                    | 1     |
| Nonanal                        |                         | 1                    |                    | 1     |
| Ethyl butanoate                |                         | 100                  |                    | 1     |
| Butyl acetate                  |                         | 70                   |                    | 1     |
| Ethyl hexanoate                |                         | 7                    |                    | 1     |
| Hexyl acetate                  |                         | 5                    |                    | 1     |
| Hexyl butanoate                |                         | 1                    |                    | 1     |
| Farnesene <sup>1</sup>         | 100                     | 4                    |                    | 1     |
| ( <i>Z</i> )-3-Hexenyl acetate | 50                      |                      | 100                | 1     |
| ( <i>Z</i> )-3-Hexenol         |                         |                      | 20                 | 1     |
| ( <i>E</i> )-2-Hexenal         |                         |                      | 3                  | 1     |
| Benzaldehyde                   |                         |                      | 20                 | 1     |
| Benzonitrile                   |                         |                      | 0.5                | 1     |
| $\beta$ -Ocimene <sup>2</sup>  | 100                     |                      |                    | 1     |

<sup>1</sup> The Australian study used (*E*)- $\beta$ -farnesene and the Chinese study used a mixture of farnesene isomers. We used a mixture of farnesene isomers

<sup>2</sup> The Australian study used (*E*)- $\beta$ -ocimene, and we used a mixture of  $\beta$ -ocimene isomers

<sup>a</sup> Il'ichev et al., 2009

<sup>b</sup> Lu et al., 2012 (JM blend, Table 2)

<sup>c</sup> Piñero and Dorn, 2007

**Table 3** Field and flight tunnel experimental details

| Exp. | Objective  | Start-end dates                      | Sex pheromone | Pheromone:host-plant ratio  |
|------|--|--------------------------------------|---------------|---|
| 1    | Do host-plant blends attract <i>G. molesta</i> ? Compare host-plant blends alone, at two doses and in two dispenser types (mineral oil in microcentrifuge tube vs <i>n</i> -hexane in rubber septum) | December 21, 2012 - January 29, 2013 | 80 $\mu$ g    | (Host-plant blends tested alone, two doses, in <i>n</i> -hexane or in mineral oil)<br>Australian: 100 mg and 10 mg<br>Chinese: 189 mg and 19 mg<br>Swiss: 143 mg and 14 mg<br>Total: 140 mg and 14 mg |
| 2    | Is there sex pheromone and host-plant synergism? Compare sex pheromone and host-plant blends at one host-plant dose.   | January 29 - February 12, 2013       | 16 $\mu$ g    | Pher.:Australian, 1: 6,250<br>Pher.:Chinese, 1: 11,812<br>Pher.:Swiss, 1: 9,062<br>Pher.:Total, 1: 8,750<br>(Host-plant alone same as high conc. of exp. 1)   |
| 3    | Is there sex pheromone and host-plant inhibition? Is inhibition dose-dependent? Compare sex pheromone and host-plant blends at several host-plant doses. Test additional host-plant compounds        | February 12 - 25, 2013               | 8 $\mu$ g     | Pher.:Chinese, 1:237.5, 1:7,875, 1:23,625<br>Pher.:Total, 1:175, 1:5,825, 1:17,500<br>Pher.: $\beta$ -Ocimene, 1:375<br>Pher.:Terpinyl acetate, 1:375   |
| 4    | Test sex pheromone and host-plant blends of field experiments in the wind tunnel   | February 8 - 27, 2013                | 1ng           | Pher.:Australian/Chinese/Swiss, 1:10, 1:100, 1:1,000, 1:10,000<br>(Plant alone: 10 $\mu$ g)   |

**Table 4** Captures of *G. molesta* in Chile between December 21<sup>st</sup> 2012 and January 29<sup>th</sup> 2013 in traps baited with one of 4 host-plant blends in two doses and either dissolved in *n*-hexane and loaded in red rubber septa, or dissolved in mineral oil and loaded in microcentrifuge tubes (Experiment 1) ~~Solvents and sex pheromone were negative and positive controls, respectively. Only 1 female was captured in the entire experiment. Different letters within the Males columns indicate significant differences among treatments (Tukey,  $p \leq 0.05$ )~~

| Stimulus                 | Captures/trap/check (mean $\pm$ SEM) |         |                                     |                 |
|--------------------------|--------------------------------------|---------|-------------------------------------|-----------------|
|                          | <i>n</i> -Hexane in rubber septum    |         | Mineral oil in microcentrifuge tube |                 |
|                          | Males                                | Females | Males                               | Females         |
| <i>n</i> -Hexane         | 0 <sup>c</sup>                       | 0       | 0 <sup>b</sup>                      | 0               |
| Sex pheromone 80 $\mu$ g | 68.00 $\pm$ 13.99 <sup>a</sup>       | 0       | 8.96 $\pm$ 3.26 <sup>a</sup>        | 0               |
| Australian 10 mg         | 0 <sup>c</sup>                       | 0       | 0 <sup>b</sup>                      | 0               |
| Chinese 19 mg            | 0.08 $\pm$ 0.08 <sup>c</sup>         | 0       | 0 <sup>b</sup>                      | 0               |
| Swiss 14 mg              | 1.46 $\pm$ 1.46 <sup>b</sup>         | 0       | 0.04 $\pm$ 0.04 <sup>b</sup>        | 0               |
| Total 14 mg              | 0 <sup>c</sup>                       | 0       | 0 <sup>b</sup>                      | 0               |
| Australian 100 mg        | 0 <sup>c</sup>                       | 0       | 0 <sup>b</sup>                      | 0               |
| Chinese 189 mg           | 0.04 $\pm$ 0.04 <sup>c</sup>         | 0       | 0.62 $\pm$ 0.62 <sup>b</sup>        | 0               |
| Swiss 143 mg             | 0 <sup>c</sup>                       | 0       | 0 <sup>b</sup>                      | 0.04 $\pm$ 0.04 |
| Total 140 mg             | 0.38 $\pm$ 0.38 <sup>c</sup>         | 0       | 0.04 $\pm$ 0.04 <sup>b</sup>        | 0               |

Solvents and sex pheromone were negative and positive controls, respectively. Only 1 female was captured in the entire experiment. Different letters within the Males columns indicate significant differences among treatments (Tukey,  $p \leq 0.05$ )

**Table 5** Captures of *G. molesta* in Chile between January 29<sup>th</sup> and February 12<sup>th</sup> 2013 in traps baited with either, one septum of one of 4 host-plant blends, a sex pheromone septum, or sex pheromone with a host-plant blend septum (Experiment 2) Stimulus dissolved in *n*-hexane, tested as a negative control. Different letters within the male column indicate significant differences among treatments (Tukey,  $p \leq 0.05$ ). Female captures by the Australian host-plant blend and *n*-hexane not significantly different

| Stimulus                 |                          | Captures/trap/check (mean $\pm$ SEM) |                 |
|--------------------------|--------------------------|--------------------------------------|-----------------|
| Septum 1                 | Septum 2                 | Males                                | Females         |
| <i>n</i> -Hexane         |                          | 0 c                                  | 0               |
| Sex pheromone 16 $\mu$ g |                          | 25.62 $\pm$ 3.27 <sup>a</sup>        | 0               |
| Australian 100 mg        |                          | 0 <sup>d</sup>                       | 0               |
| Chinese 189 mg           |                          | 0 <sup>d</sup>                       | 0               |
| Swiss 143 mg             |                          | 0 <sup>d</sup>                       | 0               |
| Total 140 mg             |                          | 0 <sup>d</sup>                       | 0               |
| Australian 100 mg        | Sex pheromone 16 $\mu$ g | 20.41 $\pm$ 3.05 <sup>b</sup>        | 0.28 $\pm$ 0.11 |
| Chinese 189 mg           | Sex pheromone 16 $\mu$ g | 10.28 $\pm$ 1.51 <sup>c</sup>        | 0               |
| Swiss 143 mg             | Sex pheromone 16 $\mu$ g | 20.66 $\pm$ 2.18 <sup>b</sup>        | 0               |
| Total 140 mg             | Sex pheromone 16 $\mu$ g | 10.53 $\pm$ 1.69 <sup>c</sup>        | 0               |

Stimulus dissolved in *n*-hexane, tested as a negative control. Different letters within the male column indicate significant differences among treatments (Tukey,  $p \leq 0.05$ ). Female captures by the Australian host-plant blend and *n*-hexane not significantly different

**Table 6** Captures of *G. molesta* in Chile between February 12<sup>th</sup> and February 25<sup>th</sup> 2013 in traps baited with one sex pheromone septum and a host-plant blend septum (Chinese or Total) at low, medium or high doses (Experiment 3) ~~Stimulus dissolved in *n*-hexane, tested as a negative control. Terpinyl acetate and  $\beta$ -ocimene (isomer mix) tested alone or with a sex pheromone septum. A trap baited with sex pheromone and an *n*-hexane septum tested the effect of septum number. Different letters within the male column indicate significant differences among treatments (Tukey,  $p \leq 0.05$ ). Female captures were not significantly different from *n*-hexane~~

| Stimulus                |                         | Captures/trap/check (mean $\pm$ SEM) |                 |
|-------------------------|-------------------------|--------------------------------------|-----------------|
| Septum 1                | Septum 2                | Males                                | Females         |
| <i>n</i> -Hexane        |                         | 0.21 $\pm$ 0.16 <sup>h</sup>         | 0.04 $\pm$ 0.04 |
| Sex pheromone 8 $\mu$ g |                         | 46.68 $\pm$ 6.35 <sup>bc</sup>       | 0               |
| Sex pheromone 8 $\mu$ g | <i>n</i> -Hexane        | 66.18 $\pm$ 8.48 <sup>a</sup>        | 0               |
| Chinese 1.9 mg          | Sex pheromone 8 $\mu$ g | 48.71 $\pm$ 9.83 <sup>b</sup>        | 0.04 $\pm$ 0.04 |
| Chinese 63 mg           | Sex pheromone 8 $\mu$ g | 39.75 $\pm$ 5.42 <sup>d</sup>        | 0               |
| Chinese 189 mg          | Sex pheromone 8 $\mu$ g | 24.57 $\pm$ 5.14 <sup>ef</sup>       | 0.04 $\pm$ 0.04 |
| Total 1.4 mg            | Sex pheromone 8 $\mu$ g | 40.71 $\pm$ 7.00 <sup>cd</sup>       | 0               |
| Total 46.6 mg           | Sex pheromone 8 $\mu$ g | 28.57 $\pm$ 4.37 <sup>e</sup>        | 0.07 $\pm$ 0.05 |
| Total 140 mg            | Sex pheromone 8 $\mu$ g | 21.46 $\pm$ 5.33 <sup>f</sup>        | 0.04 $\pm$ 0.04 |
| Terpinyl Ac. 3 mg       |                         | 0.57 $\pm$ 0.28 <sup>gh</sup>        | 0.04 $\pm$ 0.04 |
| Terpinyl Ac. 3 mg       | Sex pheromone 8 $\mu$ g | 63.82 $\pm$ 8.26 <sup>a</sup>        | 0.14 $\pm$ 0.07 |
| $\beta$ -Ocimene 3 mg   |                         | 1.46 $\pm$ 1.39 <sup>g</sup>         | 0               |
| $\beta$ -Ocimene 3 mg   | Sex pheromone 8 $\mu$ g | 52.93 $\pm$ 7.14 <sup>b</sup>        | 0.21 $\pm$ 0.08 |

Stimulus dissolved in *n*-hexane, tested as a negative control. Terpinyl acetate and  $\beta$ -ocimene (isomer mix) tested alone or with a sex pheromone septum. A trap baited with sex pheromone and an *n*-hexane septum tested the effect of septum number. Different letters within the male column indicate significant differences among treatments (Tukey,  $p \leq 0.05$ ). Female captures were not significantly different from *n*-hexane